



## ¿Alto o bajo espín? Guía para facilitar esta elección en la Teoría del Campo de los Ligantes

### *High- or low-spin complex? A guide to facilitate the selection in Ligand Field Theory*

Sara Angelica Cortes-Llamas<sup>1</sup>, José Miguel Velázquez-López<sup>1</sup>, Irma Idalia Rangel-Salas<sup>1</sup>, Morelia Eunice López-Reyes<sup>1</sup>, Alfredo Rosas-Sánchez<sup>1</sup>, Leticia Lozada-Rodríguez<sup>1</sup>, Gabriela de Jesús Soltero-Reynoso<sup>1</sup> y Saul Gallegos-Castillo<sup>1</sup>

Recepción: 01/03/2021  
Aceptación: 02/10/2021

### Resumen

La química de coordinación es un tema que resulta muy interesante para los estudiantes, debido en parte a que el razonamiento sobre el desdoblamiento de los orbitales-d puede ser confirmado con las propiedades espectroscópicas de los compuestos, particularmente sus colores. Sin embargo, en los cursos introductorios de química inorgánica es difícil para los estudiantes determinar si un complejo de coordinación homoléptico tipo Werner, tiene una configuración de alto o bajo espín. En esta contribución, proponemos dos diagramas que guían a los estudiantes para encontrar esta respuesta.

### Palabras clave

Química inorgánica; Toma de decisiones; Química de coordinación; Teoría del campo de los ligantes.

### Abstract

The study of Coordination Chemistry is a remarkably interesting topic for students, partly because the reasoning about the splitting of the *d*-orbitals can be easily confirmed with spectroscopic properties of the compounds, particularly their colors. However, in introductory inorganic chemistry courses, determining whether a homoleptic Werner-type coordination compound has a high or low-spin configuration is difficult for students. In this contribution, we propose two flowcharts that will guide the students to easily find this answer.

### Keywords

Inorganic Chemistry, Problem Solving / Decision Making, Crystal Field / Ligand Field Theory, Coordination Compounds.

<sup>1</sup> Universidad de Guadalajara, México.


The color of transition metal complexes makes Coordination Chemistry one of the most appealing topics addressed in inorganic chemistry courses. The explanation of this property in coordination compounds lies in the so-called *d*-orbital splitting. This subject becomes even more interesting when the theoretical discussion is accompanied by laboratory experiments that allow students to corroborate the nondegeneracy of *d*-orbitals due to the coordination of certain ligands to the transition metals.(Warzecha et al., 2019) However, in an introductory course on coordination chemistry, we have noticed that it becomes difficult for students to visualize whether a homoleptic Werner-type coordination compound (i.e. an octahedral or a tetrahedral complex in which all ligands bound to the metal center are identical) has a high- or a lowspin configuration; this problem arises from the fact that *d*-orbital splitting depends on several factors such as geometry, nature of ligand, oxidation state of the transition metal and the position of the metal in the periodic table (transition series). Moreover, when the coordination number (CN) is 4, it is also complicated for students to determine whether the favored geometry of the complex is tetrahedral or square planar.

Despite the fact that in the literature there is a large number of chemistry textbooks that explain outstandingly well the structure, nomenclature, molecular geometry, isomerism and bonding theories for coordination compounds,(Lawrance, 2009)-(Housecroft, Catherine E.; Sharpe, 2005)-(Miessler, Gary L.; Fischer, Paul J.; Tarr, 2014)-(House, 2008) nowadays there is no tool to systematically determine spin states in octahedral or tetrahedral complexes. To facilitate this learning process, we have designed two flowcharts (Figures 1 and 2) that assist the students to correctly analyze this *d*-orbital splitting phenomenon. These diagrams have been implemented in our introductory courses of inorganic chemistry and, over six years, we have verified that these flowcharts allow students to acquire the skills to, afterwards, deal with more challenging cases such as coordination compounds bearing non-identical ligands (heteroleptic complexes), or ligands not included in the common spectrochemical series. In our experience, these diagrams have been an excellent tool to assign the high-spin or low-spin configurations in homoleptic Werner-type compounds.

The description of the flowcharts is presented below, but first it is necessary to define important aspects related to the *d*-orbital splitting.

The flowcharts are introduced to students after the Ligand Field Theory (LFT)(Cotton, 1964)(Johnson & Graham, 2015) is discussed. According to this theory, in an octahedral geometry, ligands that generate large energy separation between *d*-orbitals (i.e., large  $\Delta_o$ ) are usually  $\pi$ -acceptors, which can perform back-donation especially with electron-rich metals. The  $\pi$ -acceptor ligands have low-energy vacant orbitals ( $\pi^*$ ) where they can favorably receive the electronic density from the metal. Some of these ligands can be easily recognized when the donor atom A presents an unsaturated bond together with another neighboring atom B (for example, M–A=B). On the other hand, ligands that generate a small  $\Delta_o$  are commonly  $\pi$ -donors, which are characterized by having at least one free electron-pair on the atom directly bound to the metal (for example, M– $\ddot{A}$ –B). These ligands can yield additional electronic density to the metallic center by forming a  $\pi$ bond. It is important to emphasize that not all ligands form  $\pi$ -interactions with the metals; for example, ammonia and amines, which are classified as  $\sigma$ -donors only. Some of these different ligands are organized in a spectrochemical series according to their ability to increase or decrease  $\Delta_o$  (Figure 3). In our experience, the description of ligands given above is useful for students to classify a ligand that is not included in the spectrochemical series.

**FIGURE 3.**  
 Spectrochemical series  
 of ligands (L = neutral  
 ligand; X = anionic  
 ligand).



It is also convenient to classify the ligands as L- and X-type ligands according to the neutral convention. L-type ligands are molecules that form a covalent-coordinated bond with the metal center; these are neutral species that formally contribute with two electrons, from a lone-pair or an unsaturation ( $\pi$ -bond), to establish the interaction with the transition metal. Examples of L-type ligands are  $\text{H}_2\text{O}$ ,  $\text{ROH}$ ,  $\text{RNH}_2$ ,  $\text{NH}_3$ ,  $\text{CO}$ , alkenes, and alkynes. The X-type ligands are anions that form a covalent bond with the metal, e.g.,  $\text{OH}^-$ ,  $\text{RO}^-$ ,  $\text{N}_3^-$ ,  $\text{CN}^-$ , and halide ions. If the anionic ligand forms more than one covalent interaction with the metal center, each interaction should be considered for the classification, for example, the oxo ligand ( $\text{O}^{2-}$ ,  $\text{M}=\text{O}$ ) is classified as an  $\text{X}_2$  ligand. Meanwhile, it is useful to determine four parameters that generally describe a metal complex. (Astruc, 2007) The first parameter is the “Number of Valence Electrons,  $NVE$ ” (Scheme 1, eq. 1), which indicates the number of total electrons around the metal center within a complex, i.e., valence electrons of the metal in its neutral state plus the electrons provided by the ligands and the electrons provided by coordination sphere charge ( $q$ ) (one more electron for each negative charge and one less electron for each positive charge). In this method, the X- and L-type ligands yield 1 and 2 electrons, respectively. The second parameter is the “Number of Non-Bonding Electrons,  $NNBE$ ” (Scheme 1, eq. 2), which determines the number of electrons that the metal has not used to form bonds. This parameter is important because it provides a quick visualization of the expected reactivity of a complex. For example, when a metal in a complex has  $NNBE = 0$ , it cannot be oxidized as it has no electrons to donate (nobody can give what they do not have!). In the literature,  $NNBE$  parameter is usually represented as  $d^{NNBE}$ . The third parameter is the “Oxidation State,  $OS$ ” (Scheme 1, eq. 3), which represents the hypothetical charge that a metal has if the M-X bonds are completely ionic (which is not true since the M-X bonds in coordination compounds are generally polar covalent); nonetheless, it is a useful parameter as a first idea about the lack or excess of electronic density on the metal. Finally, the fourth parameter is the “Coordination Number,  $CN$ ” (Scheme 1, eq. 4), which is the number of occupied positions by the ligands around the metal. The main  $CNs$  are 2, 3, 4, 5 and 6, and the most common values are 4 and 6. In Table 1, we present two examples for the calculation of these four parameters.

**SCHEME 1.** Equations to obtain  $NVE$ ,  $NNBE$ ,  $OS$  and  $CN$  parameters. ( $nM$  = number of valence electrons of the metal in its ground state;  $nL$  = number of neutral L-type ligands;  $nX$  = number of anionic X-type ligands;  $q$  = coordination sphere charge).

$$NVE = nM + 2nL + nX - q \quad \text{eq. 1}$$

$$NNBE = nM - nX - q \quad \text{eq. 2}$$

$$OS = nX + q \quad \text{eq. 3}$$

$$CN = nL + nX \quad \text{eq. 4}$$


**TABLE 1.** Determination of the four basic parameters in the description of two selected coordination complexes.

$[\text{Fe}(\text{ox})_3]^{3-}$	$[\text{Co}(\text{NH}_3)_6]^{2+}$
$[\text{Fe}(\text{X}_2)_3]^{3-} = [\text{FeX}_6]^{3-}$ ( $\text{ox}^{2-}$ = oxalate is $\text{X}_2$ -type ligand, meaning that oxalate is a bidentate ligand forming two Fe-O covalent bonds)	$[\text{CoL}_6]^{2+}$ ( $\text{NH}_3$ is L-type ligand)


$NVE = 8 + 2(0) + 6 - (-3) = 17$ $NNBE = 8 - 6 - (-3) = 5 (d^5 \text{ complex})$ $OS = 6 + (-3) = 3$ $CN = 0 + 6 = 6$	$NVE = 9 + (2)(6) + 0 - (+2) = 19$ $NNBE = 9 - 0 - (+2) = 7 (d^7 \text{ complex})$ $OS = 0 + 2 = 2$ $CN = 6 + 0 = 6$
---	--

Having defined the *NVE*, *NNBE*, *OS* and *CN* parameters and the equations for their calculation, we can now review the flowcharts in Figures 1 and 2.

**FIGURE 1.** Flowchart of the high- vs. low-spin assignment for octahedral complexes (coordination number = 6). SS = Spectrochemical Series. \*Exceptions have been observed: *OS* = +3 and  $\sigma$ -donor only chelate ligands = Low-spin state.



**FIGURE 2.** Flowchart of the high- vs. low-spin assignment for complexes with coordination number = 4. \*Some exceptions have been observed: *OS* > +3 and absence of  $\pi$ -donor ligands = Low-spin state.



We start with the coordination number of six ( $CN = 6$ ); the most common geometry in this case is octahedral, so only such geometry is considered. According to the flowchart for a complex with  $CN = 6$  (Figure 1), the considerations to take into account for the assignment of the spin configuration are as follows:

1. If the metal belongs to the first transition series, then:
  - a) If it is  $Co^{3+}$  ( $d^6$ ), a low-spin complex is formed.(Åkesson et al., 1994) In this exception, a great energy of stabilization of the crystalline field is generated when six electrons of  $Co^{3+}$  are placed in the  $t_{2g}$  sublevel compared to that obtained in a high-spin configuration.
  - b) If it is a  $M^{2+}$  or  $M^{3+}$  ion (except for  $Co^{3+}$ ) bound to  $\pi$ -donor or  $\sigma$ -donor only ligands (ending with ethylenediamine ligand, *en*, in the spectrochemical series of the Figure 3), a high-spin complex is formed). However, the combination of  $M^{3+}$  with chelate  $\sigma$ -donors only, generates low-spin complexes.(POON & ANDREW, 1979).(Comba, 1994).(Bhowmick et al., 2018).(Machida et al., 1986)<sup>a</sup> high-spin  $Ni(II)$  This high field may be due to the combination of a high oxidation state and excellent sigma donor ligands.
  - c) If it is a  $M^{2+}$  or  $M^{3+}$  ion (except for  $Co^{3+}$ ) bound to  $\pi$ -acceptor ligands, a low-spin complex is formed.
2. If the metal belongs to the second or third series of transition metals, the complex will tend to have a low-spin configuration, regardless of its oxidation state and the nature of the ligands to which it is bound.

For example, for iron, which is a metal from the first transition series, hexaaqua complexes  $[Fe(H_2O)_6]^{2+}$  ( $NVE = 18$ ;  $NNBE = 6$ ;  $OS = +2$ ;  $CN = 6$ ) and  $[Fe(H_2O)_6]^{3+}$  ( $NVE = 17$ ;  $NNBE = 5$ ;  $OS = +3$ ;  $CN = 6$ ) are high-spin (4 and 5 unpaired electrons, respectively); likewise  $[FeBr_6]^{3-}$  ( $NVE = 17$ ;  $NNBE = 5$ ;  $OS = +3$ ;  $CN = 6$ ) is a high-spin complex(Jouini et al., 1983) because bromide is a  $\pi$ -donor ligand. Meanwhile,  $[Fe(NO_2)_6]^{3-}$ ,  $[Fe(CN)_6]^{3-}$  ( $NVE = 17$ ;  $NNBE = 5$ ;  $OS = +3$ ;  $CN = 6$ ) and  $[Fe(CN)_6]^{4-}$  ( $NVE = 18$ ;  $NNBE = 6$ ;  $OS = +2$ ;  $CN = 6$ ) are low-spin complexes(Clark et al., 1954).(Koubek & Elert, 1996) (1, 1 and 0 unpaired electrons, respectively) because  $NO_2^-$  and  $CN^-$  are  $\pi$ -acceptors.

With respect to  $\sigma$ -donor only ligands, they tend to form high-spin complexes. For example,  $[M(NH_3)_6]^{2+}$  ( $M = Cr, Fe, Mn, Co$  and  $Ni$ )(Redaktion, 2006) as well as  $[M(en)_3]^{2+}$  ( $M = Mn, Co, Ni$ )(Cortijo et al., 2020) are high-spin complexes. Nevertheless, in the case of  $M^{3+}$  with amine chelate ligands, being excellent  $\sigma$ -donor ligands, they tend to generate low-spin complexes, as is the case of  $[Fe(en)_3]^{3+}$ . (Renovitch & Baker, 1968) A very illustrative example is found in the laboratory practice proposed by Morrow with 1,8-dinitro-3,6,10,13,16,19-hexaazabicyclo-(6,6,6)eicosanelso ligand called dinitrosarcophagine or “dinosaur”;(Burns et al., 2016) this hexamine ligand forms a high-spin complex with  $Co^{2+}$ , while the analogous complex with  $Co^{3+}$  is a low-spin complex.

For the second and third series of transition metals, low-spin complexes are usually observed even with  $\pi$ -donor ligands, for example  $[RuF_6]^{3-}$  ( $OS = +3$ ),(Allen et al., 1973) or  $[Ru(H_2O)_6]^{3+}$  ( $OS = +3$ ).(Bernhard et al., 1984)

The flowchart in Figure 1, facilitates the initial high- or low-spin configuration assignment in homoleptic compounds. However, we have observed that this first step allows



the students to develop the ability to assign the spin state in a wide variety of Werner-type complexes. For example, the ethylenediaminetetraacetate (EDTA) ligand is not included in the spectrochemical series of Figure 3. For this hexadentate ligand, would we expect high- or low-spin complexes according to the flowchart in Figure 1? Bearing four acetate groups ( $\pi$ -donor ligands) and two amino groups ( $\sigma$ -donor only ligands) we would expect EDTA to form high-spin complexes with  $M^{2+}$  and  $M^{3+}$  metals of the first transition series. However, in the case of Co(III) we would expect a low-spin complex. For metals of the second and third transition series, we would expect complexes with only low-spin configuration. This assignment agrees well with the reports in the literature. Indeed,  $Co^{3+}$  forms a low-spin complex with EDTA (Douglas & Radanović, 1993) while  $Mn^{2+}$  and  $Fe^{3+}$  (from the first transition series) form high-spin complexes; (Hessler et al., 1994) on the other hand, both  $Ru^{2+}$  and  $Ru^{3+}$  (from the second transition series) form low-spin complexes. (Rocha et al., 2002)

Regarding the coordination number of four ( $CN = 4$ ), there are several reports in the literature on the factors that influence the geometry and spin state preferred by a compound. (Cirera et al., 2004, 2008; Poli, 1996). For instance, an excellent analysis of tetrahedral geometry for homoleptic complexes was carried out by Alvarez et al. where they proposed the so-called “Magic Cube”. Taking these reports into account, we have extracted the information to generate a flowchart (Figure 2) to facilitate students a correct determination of geometry and spin states. According to the flowchart for a complex with  $CN = 4$ , the considerations to consider are as follows:

1. If  $NNBE = 0, 1, 2, 5$  or  $10$ , the complexes will adopt a tetrahedral geometry. In this geometry, the complexes are high spin regardless of the transition series of the metal, ligands, and oxidation state. However, some exceptions observing tetrahedral low-spin complexes have been found, especially when the metal oxidation state is  $\geq +4$ , with no  $\pi$ -donor ligands and metals mainly from the second or third transition series. (Byrne & Theopold, 1989; Hay-Motherwell et al., 1991) (Jenkins et al., 2002).
2. If the complexes have  $NNBE \neq 0, 1, 2, 5$  or  $10$  and the metal belongs to the first transition series, then:
  - a) If the metal is bound to  $\pi$ -acceptor ligands, the favored geometry is square-planar, low-spin configuration.
  - b) If the metal is bound to  $\pi$ -donor ligands, the favored geometry is tetrahedral, high-spin configuration.
3. If the complex has  $NNBE \neq 0, 1, 2, 5$  or  $10$  and the metal belongs to the second and third transition series, the square-planar geometry is favored regardless of the ligands.

Although steric effects and chelate ligands have a great influence on geometry, the guide of Figure 2 is a great help for homoleptic coordination compounds. For example, the complexes  $[MnO_4]^-$ ,  $[MnCl_4]^{2-}$  and  $[Ni(CO)_4]$ , which have  $NNBE$  of 0, 5, and 10, respectively, exhibit tetrahedral geometry regardless of the oxidation state of the metal (+7, +2 and 0, respectively) or whether the metal is bound to a  $\pi$ -donor ( $O^{2-}$  and  $Cl^-$ ) or a  $\pi$ -acceptor ligand (CO).

Another example is found in  $[NiCl_4]^{2-}$  and  $[Ni(CN)_4]^{2-}$ . (Manch, 1961) Nickel belongs to the first series of transition metals and, in both compounds,  $NNBE = 8$ . Because chlorides are  $\pi$ -donor ligands,  $[NiCl_4]^{2-}$  has a tetrahedral geometry; on the contrary, cyanides are

$\pi$ -acceptor ligands, so  $[\text{Ni}(\text{CN})_4]^{2-}$  has a square-planar geometry. In contrast, for the  $[\text{PdCl}_4]^{2-}$  and  $[\text{Pd}(\text{CN})_4]^{2-}$  complexes, (Börgel et al., 2016) where palladium is an element of the second series of transition metals, both complexes are square-planar regardless of whether the metal is bound to a  $\pi$  donor ( $\text{Cl}^-$ ) or a  $\pi$ -acceptor ( $\text{CN}^-$ ) ligand.

Due to the simplicity of these flowcharts, we consider them a simple and practical guide for students in introductory courses on Coordination Chemistry. The use of these diagrams is suggested after reviewing of the concepts of the Ligand Field Theory. Finally, it is important to emphasize that although the use of these diagrams allows the correct analysis of a wide variety of complexes, their use should be handled with caution since there are exceptions (some of them have been commented in this work).

## Acknowledgments

This work was supported by the program “Fortalecimiento a la Investigación y el Posgrado, Edición 2020 – Universidad de Guadalajara”.

## References

- Åkesson, R., Pettersson, L. G. M., Sandström, M., & Wahlgren, U. (1994). Ligand Field Effects in the Hydrated Divalent and Trivalent Metal Ions of the First and Second Transition Periods. *Journal of the American Chemical Society*, 116(19), 8691–8704. <https://doi.org/10.1021/ja00098a032>
- Allen, G. C., El-Sharkawy, G. A. M., & Warren, K. D. (1973). Electronic Spectra of Some Hexafluoro Anions of the 4d Series. *Inorganic Chemistry*, 12(10), 2231–2237. <https://doi.org/10.1021/ic50128a004>
- Astruc, D. (2007). *Organometallic Chemistry and Catalysis*. Berlin, Germany: Springer Berlin Heidelberg <https://doi.org/10.1007/978-3-540-46129-6>
- Bernhard, P., Stebler, A., & Ludi, A. (1984). Electronic Properties of Hexaaquaruthenium(III): EPR and Optical Spectra of  $\text{Ru}(\text{H}_2\text{O})_6^{3+}$  in Alum Crystals. *Inorganic Chemistry*, 23(14), 2151–2155. <https://doi.org/10.1021/ic00182a033>
- Bhowmick, I., Roehl, A. J., Neilson, J. R., Rappé, A. K., & Shores, M. P. (2018). Slow magnetic relaxation in octahedral low-spin Ni(III) complexes. *Chemical Science*, 9(31), 6564–6571. <https://doi.org/10.1039/c7sc04482h>
- Börgel, J., Campbell, M. G., & Ritter, T. (2016). Transition Metal d-Orbital Splitting Diagrams: An Updated Educational Resource for Square Planar Transition Metal Complexes. *Journal of Chemical Education*, 93(1), 118–121. <https://doi.org/10.1021/acs.jchemed.5b00542>
- Burns, P. J., Tsitovich, P. B., & Morrow, J. R. (2016). Preparation of a Cobalt(II) Cage: An Undergraduate Laboratory Experiment That Produces a ParaSHIFT Agent for Magnetic Resonance Spectroscopy. *Journal of Chemical Education*, 93(6), 1115–1119. <https://doi.org/10.1021/acs.jchemed.5b00818>

- Byrne, E. K., & Theopold, K. H. (1989). Synthesis, Characterization, and Electron-Transfer Reactivity. *Journal of the American Chemical Society*, *111*, 3887–3896.
- Cirera, J., Alemany, P., & Alvarez, S. (2004). Mapping the Stereochemistry and Symmetry of Tetracoordinate Transition-Metal Complexes. *Chemistry - A European Journal*, *10*(1), 190–207. <https://doi.org/10.1002/chem.200305074>
- Cirera, J., Ruiz, E., & Alvarez, S. (2008). Stereochemistry and spin state in four-coordinate transition metal compounds. *Inorganic Chemistry*, *47*(7), 28 |71-2889. <https://doi.org/10.1021/ic702276k>
- Clark, H. C., Curtis, N. F., & Odell, A. L. (1954). Exchange reactions and magnetic susceptibilities of complex salts. Part I. Some complex salts of ferric iron. *Journal of the Chemical Society (Resumed)*, 63–66. <https://doi.org/10.1039/JR9540000063>
- Comba, P. (1994). Prediction and Interpretation of EPR Spectra of Low-Spin Iron(III) Complexes with the MM-AOM Method. *Inorganic Chemistry*, *33*(20), 4577–4583. <https://doi.org/10.1021/ic00098a027>
- Cortijo, M., Valentín-Pérez, Á., Rouzières, M., Clérac, R., Rosa, P., & Hillard, E. A. (2020). Tris(ethylenediamine) Cobalt(II) and Manganese(II) Nitrates. *Crystals*, *10*(6), 472. <https://doi.org/10.3390/cryst10060472>
- Cotton, F. A. (1964). Ligand field theory. *Journal of Chemical Education*, *41*(9), 466–476. <https://doi.org/10.1021/ed041p466>
- Douglas, B. E., & Radanović, D. J. (1993). Coordination chemistry of hexadentate EDTA-type ligands with M(III) ions. *Coordination Chemistry Reviews*, *128*(1–2), 139–165. [https://doi.org/10.1016/0010-8545\(93\)80027-3](https://doi.org/10.1016/0010-8545(93)80027-3)
- Hay-Motherwell, R. S., Wilkinson, G., Hussain-Bates, B., & Hursthouse, M. B. (1991). Synthesis and structure of iridium tetramesityl. *Polyhedron*, *10*(12), 1457–1458. [https://doi.org/10.1016/S0277-5387\(00\)81284-X](https://doi.org/10.1016/S0277-5387(00)81284-X)
- Hessler, D. P., Frimmel, F. H., Oliveros, E., & Braun, A. M. (1994). Solvent Isotope Effect on the Rate Constants of Singlet-Oxygen Quenching by edta and Its Metal Complexes. *Helvetica Chimica Acta*, *77*(3), 859–868. <https://doi.org/10.1002/hlca.19940770325>
- House, J. E. (2008). Chemistry of Coordination Compounds. In *Chemistry of Coordination Compounds*. Vermont, USA: Elsevier.
- Housecroft, Catherine E.; Sharpe, A. G. (2005). Inorganic chemistry. In *Inorganic Chemistry*. Chelmsford, Great Britain: Prentice-Hall.
- Jenkins, D. M., Di Bilio, A. J., Allen, M. J., Betley, T. A., & Peters, J. C. (2002). Elucidation of a low spin cobalt(II) system in a distorted tetrahedral geometry. *Journal of the American Chemical Society*, *124*(51), 15336–15350. <https://doi.org/10.1021/ja026433e>
- Johnson, B. J., & Graham, K. J. (2015). A Guided Inquiry Activity for Teaching Ligand Field Theory. *Journal of Chemical Education*, *92*(8), 1369–1372. <https://doi.org/10.1021/acs.jchemed.5b00019>



- Jouini, N., Guen, L., & Tournoux, M. (1983). Structure cristalline et proprietes magnetiques de la variete basse temperature  $TlFeBr_3$ . *Materials Research Bulletin*, 18(4), 491–497. [https://doi.org/10.1016/0025-5408\(83\)90141-1](https://doi.org/10.1016/0025-5408(83)90141-1)
- Koubek, E., & Elert, M. (1996). A demonstration of crystal-field effects in octahedral complexes. *Journal of Chemical Education*, 73(10), 947. <https://doi.org/10.1021/ed073p947>
- Lawrance, G. A. (2009). Introduction to Coordination Chemistry. In *Introduction to Coordination Chemistry*. Wiltshire, Great Britain: Wiley <https://doi.org/10.1002/9780470687123>
- Machida, R., Kimura, E., & Kushi, Y. (1986). Coordination of Pentadentate Macrocycles, Doubly Deprotonated 1,4,7,10,13-Pentaazacyclohexadecane-14,16-dione, to High-Spin Nickel(II) and Low-Spin Nickel(III). X-ray Study of a Novel Redox System. *Inorganic Chemistry*, 25(19), 3461–3466. <https://doi.org/10.1021/ic00239a029>
- Manch, W. (1961). The structure and spectra of nickel (II) and copper(II) complexes. *Journal of Chemical Education*, 38(4), 192–201. <https://doi.org/10.1021/ed038p192>
- Miessler, Gary L.; Fischer, Paul J.; Tarr, D. A. (2014). Inorganic Chemistry. In *Inorganic Chemistry*. New Jersey, USA: Prentice-Hall, <https://doi.org/9781292020754>
- Poli, R. (1996). Open-shell organometallics as a bridge between Werner-type and low-valent organometallic complexes. The effect of the spin state on the stability, reactivity, and structure. *Chemical Reviews*, 96(6), 2135–2204. <https://doi.org/10.1021/cr9500343>
- Poon, C. K., Andrew, M. W. (1979). Structural and Mechanistic Studies of Coordination Compounds. 23 Evidence for Dissociative Substitution Reactions of Some trans-Acido(isothiocyanato)(1,4,8,11-tetraazacyclotetradecane)iron(III) Complexes, trans-[Fe(cyclam)(NCS)X]<sup>+</sup>, X<sup>-</sup> = NCS<sup>-</sup>, CH<sub>2</sub>ClCOO<sup>-</sup>, C. *Inorganic Chemistry*, 18(5), 1277–1283.
- Redaktion, D. (2006). Zeitschrift fur Anorganische und Allgemeine Chemie: Editorial. *Zeitschrift Fur Anorganische Und Allgemeine Chemie*, 632(12–13), 1917. <https://doi.org/10.1002/zaac.200690020>
- Renovitch, G. A., & Baker, W. A. (1968). Magnetic and spectral properties of [Fe(en)<sub>3</sub>]Cl<sub>3</sub>. *Journal of the American Chemical Society*, 90(13), 3585–3587. <https://doi.org/10.1021/ja01015a064>
- Rocha, R. C., Rein, F. N., & Toma, H. E. (2002). Electrochemical parametrization of a series of penta- and tetradentate EDTA complexes of ruthenium. *Inorganic Chemistry Communications*, 5(10), 891–896. [https://doi.org/10.1016/S1387-7003\(02\)00599-3](https://doi.org/10.1016/S1387-7003(02)00599-3)
- Warzecha, E., Berto, T. C., Wilkinson, C. C., & Berry, J. F. (2019). Rhodium Rainbow: A Colorful Laboratory Experiment Highlighting Ligand Field Effects of Dirhodium Tetraacetate. *Journal of Chemical Education*, 96(3), 571–576. <https://doi.org/10.1021/acs.jchemed.6b00648>